

DST-206 Pharyngitis (Sore Throat): Pediatric

DEFINITION

A painful condition of the oropharynx associated with infection of the mucus membranes of the pharynx and the palatine tonsils. The peak prevalence is found in children less than 5 years.

Nurses with Remote Practice Certified Practice designation (RN(C)s) are able to treat children with pharyngitis who are **1 year of age and older**.

Potential Causes

Infectious

Viruses

- Adenovirus
- Parainfluenza virus
- Epstein –Barr
- Coxsackievirus
- Herpes simplex virus
- Enterovirus (more common in children less than 3 years of age)
- Influenza virus

Bacterial

- Group A beta-haemolytic strep (GAS) (*streptococcus pyogenes*)
- *Mycoplasma pneumoniae* (10% of adolescents)
- *Neisseria gonorrhoeae* or *Chlamydia trachomatis* (related to sexual activity)
- Chlamydia pneumoniae
- Diphtheriae

Non-infectious

- Allergic rhinitis
- Sinusitis with post nasal drip
- Mouth breathing
- Trauma
- GERD (gastroesophageal reflux disease)

PREDISPOSING RISK FACTORS

- Previous episodes of pharyngitis or tonsillitis
- Smoking, exposure to cigarette smoke
- Overcrowding
- Immunocompromised
- Steroids, oral or inhaled
- Diabetes mellitus
- Oral sex

TYPICAL FINDINGS OF SORE THROAT

See Appendix 1 for pathogens and clinical appearance of tonsils

Note: Always consider the potential for epiglottitis and airway obstruction. If symptoms of airway distress, tripoding, stridor, dysphagia, drooling and anxiety exist, do not exam the child's mouth or throat, but immediately consult with or refer the client to a physician or nurse practitioner.

Bacterial

History

- Acute onset
- Very sore throat
- Absence of cough and coryza
- Fever
- Headache
- May have nausea, vomiting, abdominal pain
- General malaise

Physical Assessment

- Significant fever
- Tachycardia
- Weigh until 12 years of age for medication calculations
- Pharyngeal and tonsillar erythema
- Petechiae of soft palate
- Tonsillar exudate (particularly with streptococcal infection, diphtheria or mononucleosis)
- Anterior cervical lymphadenopathy
- Erythematous "sandpaper" rash of scarlet fever (may be present with streptococcal infection)
- Erythematous rash (particularly if child is receiving amoxicillin)
- Lymphadenopathy with splenic enlargement in children with mononucleosis
- Koplik spots

Viral

History

- Acute sore throat combined with symptoms consistent with a viral upper respiratory tract infection (rhinorrhea, cough and often hoarseness)

Physical Assessment

- Fever (low-grade to significant)
- Tachycardia
- Weigh until 12 years of age for medication calculations
- Pharyngeal and tonsillar erythema and swelling
- Petechiae of soft palate
- Tonsillar exudate similar to that occurring with bacterial infection may be present, particularly in adenovirus pharyngotonsillitis
- Anterior cervical lymphadenopathy
- Vesicles and ulcers may be present with coxsackievirus infection or herpes
- Hepato- and splenomegaly

Note: It is often impossible to distinguish clinically between bacterial and viral pharyngitis. Most pharyngitis is due to viruses (up to 70% in the pediatric population) and does not require treatment with antibiotics. For this reason, it is important to utilize a sore throat score and diagnostic testing as available.

Criteria		Points
Temperature > 38° Celsius		1
Absence of cough		1
Swollen, tender anterior cervical nodes		1
Tonsillar swelling or exudates		1
Age 3-14 years		1
Age 15-44 years		0
Total Score	Risk of Streptococcal infection (%)	Suggested Management
0 to 1	1-10 %	No culture or antibiotic required
2-3	11-35%	Perform culture or rapid strep test. Treat only if test is +
4 or more	51-53%	Start antibiotic therapy if patient situation warrants (e.g., high fever or clinically unwell) If culture or rapid strep test performed and negative, discontinue antibiotic

Note: Treatment with antibiotics may be warranted regardless of the score if there is a concern such as:

- household contact with streptococcal infection,
- a community epidemic of streptococcal infection,
- a history of rheumatic fever, valvular heart disease, or immunosuppression, or
- a population in which rheumatic fever remains a problem

DIAGNOSTIC TESTS

- Rapid strep test (if available)
- Throat swab for culture and sensitivity
- If the child is greater than 2 years old, culture the throat before treatment or do rapid *Strep* antigen test (if available); if negative, do throat culture.
- Monospot if suspect viral

MANAGEMENT AND INTERVENTIONS

Bacterial

Goals of Treatment

- Control pain and fever
- Prevent complications
- Rapid reduction in infectivity
- Prevent spread of Group A Streptococcus
- Decrease antibiotic resistance

Non-Pharmacologic Interventions

- Rest and increase fluid intake
- Avoidance of irritants (smoke)

- Saline gargles (1tsp of salt in 1 cup of warm water)
- Increase room humidity

Pharmacologic Interventions

Note: All doses must be calculated by weight up until age 12. Pediatric doses should not exceed recommended adult doses.

- To relieve pain:
 - acetaminophen 10-15 mg/kg, po q4-6h prn. Maximum dose 75mg/kg/24hr, or a total of 4,000mg/24hr, whichever is less, or
 - ibuprofen 5-10mg/kg, po q6-8h prn. Maximum dose 40mg/kg/24hr
- Pen VK 40mg/kg/day divided bid for 10 days. Amoxicillin is no longer recommended
- In case of unavailability of the previously listed antibiotics, or allergies to the above antibiotics:
 - Cephalexin 40 mg/kg/24hr divided bid for 10 days. (DO NOT USE if patient has a severe anaphylactic reaction to penicillin.)

OR

- Azithromycin 20 mg/kg po daily for 3 days (Maximum dose 500 mg/day)

Pregnant and Breastfeeding Women

- Acetaminophen, penicillin VK, amoxicillin, azithromycin and cephalexin may be used as listed above.
- DO NOT USE ibuprofen in pregnant patients.
- **If the infection has been determined to be due to chlamydia or gonorrhea, please refer to the appropriate STI DST.**

Viral

Goals of treatment

- Relieve symptoms
- Supportive care

Non-pharmacological Interventions

- Rest
- Increase oral fluids
- Avoid irritants
- Warm saline gargles qid (1 tsp. of salt in 1 cup of warm water)

Pharmacological Interventions

Note: All doses must be calculated by weight up until age 12. Pediatric doses should not exceed recommended adult doses.

- **To relieve pain:**
 - acetaminophen 10-15 mg/kg, po q4-6h prn. Maximum dose 75mg/kg/24hr or a total of 4,000mg/24hr, whichever is less
- OR**
- ibuprofen 5-10mg/kg, po q6-8h prn. Maximum dose 40mg/kg/24hr

Pregnant and Breastfeeding Youth

- Acetaminophen may be used in pregnant and breastfeeding youth
- DO NOT use ibuprofen in pregnancy patients.

POTENTIAL COMPLICATIONS

- Rheumatic fever (group A strep)
- Acute Glomerulonephritis (group A strep)
- Peritonsillar abscess
- Epiglottitis
- **Retropharyngeal abscess**
- Otitis media
- Sinusitis
- Splenomegaly (Epstein Barr Virus or Infectious Mononucleosis)
- Pneumonia
- Bacterial meningitis

CLIENT/CAREGIVER EDUCATION AND DISCHARGE INFORMATION

- Advise on condition, timeline of treatment and expected course of disease process
- Saline gargles as described above
- Counsel parents/caregiver about appropriate use of medication (dosage, compliance, follow-up)
- If child has any difficulty swallowing, seek help immediately

MONITORING AND FOLLOW-UP

- Return to clinic in 48 hours if awaiting culture results
- Return for care if no improvement in 48 hours

CONSULTATION AND/OR REFERRAL

- Consult a physician or nurse practitioner if child has recurrent bouts of GAS pharyngitis/tonsillitis: greater than 5 episodes in one year.

DOCUMENTATION

- As per agency policy

REFERENCES

More recent editions of any of the items in the Reference List may have been published since this DST was published. If you have a newer version, please use it.

Anti-Infective Review Panel. (2012). *Anti-infective guidelines for community-acquired infections*. Toronto, ON: MUMS Guideline Clearinghouse.

Blondel-Hill, E., & Fryters, S. (2012). *Bugs and drugs: An antimicrobial infectious diseases reference*. Edmonton, AB: Alberta Health Services.

Campisi, P., & Tewfik, T. L. (2003). [Tonsillitis and its complications](#). *Canadian Journal of Diagnosis*, 20(3), 99-105.

Canadian Pharmacists Association. (2014). *Therapeutic choices* (7th ed.). Ottawa, ON: Author.

Cash, J. C., & Glass, C. A. (Eds.). (2014). *Family practice guidelines* (3rd ed.). New York, NY: Springer.

Chen, Y. A., & Tran, C. (Eds.). (2011). *The Toronto notes 2011: Comprehensive medical reference and review for the Medical Council of Canada Qualifying Exam Part 1 and the United States Medical Licensing Exam Step 2* (27th ed.). Toronto, ON: Toronto Notes for Medical Students.

Chiappini, E., Principi, N., Mansi, N., Serra, A., De Masi, S., Camaioni, A., ... de Martino, M. (2012). Management of acute pharyngitis in children: Summary of the Italian National Institute of Health guidelines. *Clinical Therapeutics*, 34(6), 1442-1458.

DynaMed. (2016, March 25). [Antibiotics for streptococcal pharyngitis](#).

DynaMed. (2016, March 25). [Pharyngitis](#).

DynaMed. (2016, March 25). [Streptococcal pharyngitis](#).

Esau, R. (Ed.). (2012). *British Columbia's Children's Hospital pediatric drug dosage guidelines* (6th ed.). Vancouver, BC: Children's & Women's Health Centre of B.C.

Gore, J. M. (2013). Acute pharyngitis. *JAAPA: Journal of the American Academy of Physician Assistants*, 26(2), 57-58.

Hersh, A. L., Jackson, M. A., & Hicks, L. A. (2013). Principles of judicious antibiotic prescribing for upper respiratory tract infections in pediatrics. *Pediatrics*, 132(6), 1146-1154.

Jensen, B., & Regier, L. D. (Eds.). (2014). *RxFiles: Drug comparison charts* (10th ed.). Saskatoon, SK: RxFiles.

Lexicomp. (2017). Amoxicillin. Lexicomp Online.

Michigan Quality Improvement Consortium. (2013, January). *Guideline: Acute pharyngitis in children 2-18 years old* (Rev.).

National Institute for Health and Clinical Excellence (NICE). (2008, July). *Respiratory tract infections – antibiotic prescribing: Prescribing of antibiotics for self-limiting respiratory tract infections in adults and children in primary care*. Manchester, UK: Author.

Nicoteri, J. A. L. (2013). Adolescent pharyngitis: A common complaint with potentially lethal complications. *Journal for Nurse Practitioners*, 9(5), 295-300.

Shah, U. K. (2015, July 9). [Tonsillitis and peritonsillar abscess treatment & management](#).

Shulman, S.T., Bisno, A.L., Clegg, H.W., Gerber, M.A., Kaplan, E.L., Lee, G.,...Van Beneden, C. (2012). [Clinical practice guideline for the diagnosis and management of group A streptococcal pharyngitis: 2012 update by the Infectious Diseases Society of America](#). *Clinical Infectious Diseases*. 55(10), e86-e102.

University of Michigan Health System. (2013, May). [Pharyngitis. Guidelines for clinical care: Ambulatory](#) (Rev.). Ann Arbor, MI: Author.

APPENDIX 1

Table 1 – Review of pathogens that cause tonsillitis		
Pathogens		Clinical Appearance of Tonsils
Viruses	Rhinovirus, adenovirus, influenza virus, parainfluenza virus, etc.	Enlarged, erythematous.
	Coxsackie virus (herpangina).	Aphthous-like ulcers on tonsillar pillars
	Epstein-Barr virus (mononucleosis syndrome).	Very large, swollen, and dirty-grey appearance
Bacteria	Streptococcus pyogenes and other streptococcal species.	Enlarged, erythematous, with yellowish-white spots. May have membrane or purulent exudate.
Aerobic	Neisseria gonorrhoeae.	Acute purulent exudates.
	Corynebacterium diphtheriae	Exudative pharyngotonsillitis with thick pharyngeal membrane.
Anaerobic	Bacteroides species	Enlarged, erythematous.
Yeast	Candida species	White plaques with raw undersurface.
Spirochetes	Treponema pallidum (syphilis)	Oral chancres of the lip, tongue, tonsil and palate. Superficial greyish patches of mucus membrane with reddish border.
	Spirochaete denticolata and treponema vincentii (Vincent's angina)	Membrane on tonsil with underlying ulcer

Source: Campisi and Tewfik (2003) *Tonsillitis and Its Complications*